Gibbs Free Energies
C:\Courses\320\in-class\04-GibbsE.wpd; August 23, 2003 (11:20am)

formula	mineral/phase) G° _f kJ/mol
O_2	oxygen	0
Al	aluminum	0
H ₂ O	water	-237.141
Al_2O_3	corundum	-1582.228
AIO(OH)	diaspore	-918.4
AI(OH) ₃	gibbsite	-1154.889

The graph below provides a convenient way to compare some Gibbs Free Energies in the chemical system Al_2O_3 H_2O Al_2O_3 - H_2O . -100 1. Put tic marks on the -200 top line where the compositions of -300 corundum, diaspore and -400 water plot. Then move Gibbs down and put dots -500 (clearly labeled) to show Energy -600 the) G°, for each phase. of Be sure to plot very Formation -700 carefully! (kJ/mol)-800 2. Draw a line from -900 corundum to water. If -1000 diaspore plots below this line, it means that -1100 diaspore is more stable -1200 than separate water and corundum. Is it? (Note -1300 that we cannot do the -1400 same graphical analysis -1500 for the stability of gibbsite because the reaction of -1600 corundum+water to

gibbsite does not have the same number of moles on each side.)

3. Another way to compare relative stabilities is to consider the Gibbs energies of reactions. Consider the reactions:

<u>Balance</u> these reactions and use the Gibbs energy values from the table to calculate) G°_{rxn} at STP (standard temperature and pressure). Do these values come out to be less than 0? If so, diaspore and gibbsite are more stable than separate corundum + water at STP.

4. In the Al_2O_3 - H_2O system, there are two other possible chemical reactions involving three phases. List and balance them. Calculate **)** G°_{rxn} for each. For each, which is the stable side of the reaction at STP?

5. If a rock that is 100% Al₂O₃, then (considering only the Al₂O₃-H₂O system) it must contain 100% corundum. (There is no combination involving diaspore, gibbsite or water that equals 100% Al₂O₃.)

Suppose, however, you have a rock that is $50\% \text{ Al}_2\text{O}_3$ and $50\% \text{ H}_2\text{O}$. You could have any of the following three assemblages: (1) only diaspore, (2) gibbsite + corundum, (3) water + corundum.

Which of these assemblages is most stable at STP? Explain how you concluded this?