

**Problem Set #1: Geothermal gradients**

The expressions below can be used to calculate a conductive geothermal gradient for the lithosphere.

$$T(z) = Qz/K + A_0 z(b - z/2)/K + T_s \quad z < b$$

$$T(z) = Qz/K + A_0 b^2/(2K) + T_s \quad b \leq z \leq L \quad \text{where } L=100 \text{ km}$$

typical values

|        |   |     |
|--------|---|-----|
| where: | $T_s$ = surface temperature (°C)                              | 15  |
|        | $Q$ = mantle heat flow (mW/m <sup>2</sup> )                   | 30  |
|        | $K$ = thermal conductivity (W/m/deg)                          | 2.5 |
|        | $A_0 = \rho H_s$ = heat production ( $\mu$ W/m <sup>3</sup> ) | 2.0 |
|        | $b$ = characteristic depth of $A_0$ (km)                      | 10  |
|        | $z$ = depth (km)  |     |

Using a spreadsheet (e.g. Excel), plot temperature (°C) vs. depth (km) for

- the entire lithosphere (100 km), and
- the upper 35 km. Plot depth as the y-axis and “negative” (i.e. going down the page from 0 km).

**Answer the following questions:**

1. What is the temperature at the base of the lithosphere? \_\_\_\_\_ at 35 km? \_\_\_\_\_.
2. Play around with some of the parameters. What do you need to do to get 700°C at 35 km? Is there a unique solution? Which parameters do you think we know best? the least?
3. The equations given above assume that heat flow in the lithosphere is by conduction only. Is this a reasonable assumption? Why or why not?